

Comment on “On the Origin of the Highest Energy Cosmic Rays”

L. N. Epele and E. Roulet

Departamento de Física, Universidad Nacional de La Plata, CC67 (1900) La Plata, Argentina

In the paper “On the origin of the highest energy cosmic rays” [1], Stecker has analysed the implications of new estimates of the extragalactic infrared (IR) background [2] on the attenuation of the ultra-high energy cosmic ray (UHCR) nuclei, trying in particular to find an explanation for the highest energy events observed, with $E \simeq 2\text{--}3 \times 10^{20}$ eV. The result obtained was that the reduction by an order of magnitude of the IR background density with respect to the original estimates adopted in the study of these processes [3] implied essentially an increase by an order of magnitude in the attenuation length of UHCR nuclei. It was then concluded that “the highest energy CR induced air-showers could have been produced by UHCR nuclei propagating from a distance of the order of 100 Mpc!”, significantly enlarging the volume where possible UHCR sources could be located.

We want to point out here that the estimate of this effect is not appropriate, and that CR nuclei with $E \gtrsim 2 \times 10^{20}$ eV cannot reach the Earth from distances beyond ~ 10 Mpc. Indeed, ref. [1] just adapted the results of ref. [3] on the attenuation of the UHCR spectrum, assigning to the spectrum of CR nuclei coming from sources at distances of 100 and 300 Mpc (corresponding to travel times of $\sim 10^{16}$ and 3×10^{16} s) those previously obtained for travel times of 10^{15} and 3×10^{15} s. This would have been correct if the dominant opacity came from the interactions with the IR photons. However, for energies above $\sim 10^{20}$ eV the main opacity source arises from photodisintegrations with photons of the cosmic microwave background (CMB), which density is well established, rather than with the IR ones. Hence, the attenuation changes in a less pronounced way than what was assumed in ref. [1].




FIG. 1. Effective energy loss time for photodisintegration of nuclei from microwave photons (CMB), from infrared photons (IR) and the total one (solid line).

We show in figure 1 the effective energy loss time for photodisintegration from microwave photons and from

IR ones, assuming that the IR background density is a factor 10 smaller than the ‘High IR’ (HIR) flux of ref. [3], corresponding to the assumption made in ref. [1]. It is apparent that the scattering from CMB photons cannot be neglected above $E \sim 10^{20}$ eV.

In figure 2 we show the attenuation for an E^{-3} input spectrum of Fe nuclei, computed as in ref. [3]. The dashed lines are for the HIR flux of ref. [3] while the solid lines are for an IR background density smaller by an order of magnitude*. Curves from left to right correspond to travel times of 2×10^{17} , 3×10^{16} , 10^{16} , 3×10^{15} and 10^{15} s. The scaling adopted in ref. [1] is not observed, since for $E > 8 \times 10^{19}$ eV it is the CMB photons which provide the main opacity source. For sources at distances of 10 Mpc, the cutoff energy is already below 2×10^{20} eV, and it is below 10^{20} eV for distances ~ 100 Mpc.




FIG. 2. Attenuated E^{-3} differential spectrum for ultrahigh energy Fe nuclei for propagation times, from left to right, of 2×10^{17} , 3×10^{16} , 10^{16} , 3×10^{15} and 10^{15} s. Dashed lines are for the HIR density, solid lines for a density 10 times smaller.

Work partially supported by CONICET, Argentina.

-
- [1] F. W. Stecker, Phys. Rev. Letters **80**, 1816 (1998).
 - [2] M. A. Malkan and F. W. Stecker, Astrophys. J. **496**, 13 (1998).
 - [3] J.L. Puget, F.W. Stecker and J.H. Bredekamp, Astrophys. J. **205**, 638 (1976).
-

*We have not included in these figures the effect of energy losses due to pair creation, which should further suppress (slightly) the end of the spectra for travel times $\sim 3 \times 10^{16}$ s. Also not shown are the small tails due to secondary nucleons.